Sourcebook of Criminal Justice Statistics 1996, page 192

Table 2.101

Respondents' perceptions of the harmfulness of selected drug use behaviors

By age group, United States, 1994

Question: "How much do you think people risk harming themselves physically and in other ways, when they do each of the following activities?"

(Percent responding "great risk")

	Total	Age group			
	all	12 to 17	18 to 25	26 to 34	35 years
Risk behavior	ages	years	years	years	and older
<u>Marijuana</u>					
Smoke once or twice	35.6%	31.3%	21.0%	23.1%	43.4%
Smoke occasionally	39.8	32.4	24.0	28.9	48.0
Smoke regularly	59.0	57.8	46.4	49.5	65.0
Cocaine					
Try once or twice	69.8	51.7	60.5	60.6	77.8
Use occasionally	79.5	62.8	73.3	75.7	85.0
Use regularly	91.0	83.3	88.8	90.8	92.9
Use "crack" occasionally	86.3	65.2	82.1	87.6	90.6
PCP					
Try once or twice					
Use regularly	69.8	44.8	55.1	66.0	78.7
oco rogalaliy	84.5	69.5	77.8	85.2	88.5
Heroin					
Try once or twice	77.0	49.4	66.5	75.3	84.8
Use regularly	91.4	77.9	87.5	92.8	94.3
Anabolic steroids					
Use occasionally	68.9	51.2	56.6	65.7	75.8
Use regularly	86.2	78.1	81.5	85.5	88.9
,					
Alcohol					
One or two drinks nearly					
every day	28.3	28.3	25.4	26.1	29.7
Four or five drinks nearly					
every day	71.8	64.9	65.1	71.7	74.6
Five or more drinks once	50.0	50.7	40.0	54.0	04.5
or twice a week	59.0	53.7	48.3	51.9	64.5
Cigarettes					
Smoke one or more					
packs per day	66.0	51.6	60.2	65.6	69.9

Note: These data are from the 1994 National Household Survey on Drug Abuse sponsored by the U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration. Households were randomly sampled from all households in the United States and interviews were conducted throughout the year. In 1994, a split sample design was employed in order to implement a new survey questionnaire. These data are based on 17,809 completed interviews using the new questionnaire.

based on 17,809 completed interviews using the new questionnaire.

Answer alternatives to the question were "no risk," "slight risk," "moderate risk," and "great risk."

For survey methodology, see Appendix 8.

Source: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, *National Household Survey on Drug Abuse: Main Findings 1994* (Rockville, MD: U.S. Department of Health and Human Services, 1996), p. 140. Table adapted by SOURCEBOOK staff.